

Comparing different ways of horizontal information transmission on cooperation Dynamics over different social networks

Ivette C. Martínez
Universidad Simón Bolívar
Grupo de Inteligencia Artificial
Caracas, Venezuela
martinez@gia.usb.ve

Klaus Jaffe
Universidad Simón Bolívar
Laboratorio de Comportamiento
Caracas, Venezuela
kjaffe@usb.ve

ABSTRACT

An Agent Based Model was used to explore the effects of spatial social networks and of different means of horizontal information transmission over cooperation when groups provide protection against predation. We tested two ways to calculate transition probabilities governing the information diffusion of the majority's opinion: using fixed rates and using a rate proportional to group's sizes. Our results show that spatial structures affect the cooperation dynamics. Particularly in Small World Networks, cooperation is more sensible to information transmission. The type of horizontal information transmission is less important as long as over 50% of individuals follow the majority rule.

Categories and Subject Descriptors

I.6 [Simulation and Modeling]: Applications

General Terms

Design, Experimentation

Keywords

cooperation, selfish herd, complex networks, spatial effects, horizontal information transmission

1. INTRODUCTION

Biologists, economists, computer scientists and physicists have all worked to further our understanding of human and animal cooperation. Yet different premises underlay these efforts. The main difference among them is the assumption that social behavior arrived through biological evolution among animals, and that culture and rational decision making is a principal driver of the evolution of cooperation and sociality among humans [17]. Human cooperation seems to be molded by both, cultural and biological forces

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

GECCO'08 July 12–16, 2008, Atlanta, Georgia, USA.
Copyright 2007 ACM 987-1-60558-130-9/08/07 ...\$5.00.

[9]. Using theories for biological evolution has provided a fertile ground to study the dynamics of processes governed by cultural evolution, such as human cooperation [6] and economics [11]

There exist important differences between the dynamics of cultural evolution [17] and biological evolution [13]. Although both processes are often mixed up and lumped together when studying the evolution of cooperation, as done in [14]. One important feature differentiating systems driven by biological (BE) and cultural evolution (CE) is the direction of information's transmission. The transmission of information in BE is vertical (heredity), and that in CE is horizontal (imitation of behavior). This feature affects the pattern and the speed of information transmission, and is sufficient to explain important differences in the dynamics between both types of evolution [8].

Several mechanisms have been proposed to explain the emergence and maintenance of cooperation in biological terms. Hamilton [4] explains cooperation between relatives through "kin selection"; in which donor and recipient of a cooperative action are genetic relatives. Between the mechanism that have been proposed to explain cooperation between unrelated individuals we have: Direct reciprocity [18] and indirect reciprocity [12].

The study of the effects of spatiality over cooperation was introduced by Nowak and May [15]. They showed how cooperation could emerge in a population of strategies without memory when individual's relations conform a spatial structure. After Nowak and May's work, cooperation by individuals occupying spatial positions in lattices or networks that interact with their neighbors have been studied for the prisoner dilemma by several authors [7],[16]. They showed that structured populations help cooperation to evolve and maintain under certain conditions.

In this work we want to explore the effects of spatiality and the intensity of the horizontal information transmission over cooperation dynamics. To do so, we modify a one-dimensional spatial model [2] to study the differences between the dynamics of cooperative, group-forming individuals subject to a selective pressure (in this case predation). We based our model on the well known "selfish herd" concept [5] and assume that cultural and biological dynamics are driven by natural selection of the phenotypes.

2. THE MODEL

We construct an agent-based model for the study of coop-

eration dynamics based on Hamilton’s “selfish herd” [5]. The model simulates a population of interacting individuals with different social roles and different information transmission directions in environments with different spatial structure.

This model was initially proposed by Cipriani and Jaffe [2] using a cellular automaton. In this formulation, the “selfish herd” is implemented by means of the importance of group formation. In particular, group formation provides protection against predation.

Spatial structures (that represent spatial relations or contact between individuals) are modeled using graphs. The vertexes of these graphs can be occupied by individuals of any species or be empty. The neighbors of an individual in the graph form the group of this individual. These graphs are initially empty and are created at the beginning of the simulations as shown in Algorithm 1. The structure of the graph remains fixed throughout the simulation execution.

Algorithm 1 Main simulation cycle

```

1: createWorld(worldSize)
2: populateWorld()
3: for  $t = 0$  to numIterations do
4:   for each agent do
5:     agentStep()
6:   end for
7:   reapDeadAgents()
8:   repopulateWorld()
9: end for

```

The majority rule is implemented here to simulate the horizontal information transmission (H). It assumes that individuals had a given probability pT of imitating the behavior of their neighbors. Each agent executes this rule each time step, this is done by the *agentStep* algorithm, Algorithm 2. In this scenario re-population (3) is done using the initial proportion of each individual’s kind.

Algorithm 2 agentStep

```

1: neigh  $\leftarrow$  getMyNeighbours()
2: nNeighCoops  $\leftarrow$  countCooperators(neigh)
3: dead  $\leftarrow$  predate(nNeighCoops)
4: if not dead then
5:   naturalMortality()
6: end if
7: if not dead and CT then
8:   applyMajorityRule()
9: end if

```

Algorithm 3 World Repopulate

```

1:  $p_{co}$   $\leftarrow$  initial proportion of cooperators
2:  $p_{nCo}$   $\leftarrow$  initial proportion of non cooperators
3: for each empty node do
4:   create a new agent on node, cooperator with probability  $p_{co}$  or non cooperator with probability  $p_{nCo}$ 
5: end for

```

3. EXPERIMENTS

We made the implementation of the proposed model in C++. This implementation allows structured environment

for populations, in the form of complex networks, implemented as bi-bidirectional graphs. Simulations were done over populations of 10^4 individuals during 10^2 iterations.

At the beginning of simulations the graphs’ vertexes were populated by cooperators and non-cooperator, using a probability of 0.5 for each. For non-cooperators and alone cooperators the rates for predation were $pNC_o = 0.8$ and $pCo0n = 0.8$. For cooperators in groups the predation rate was $pGC_o = 0.2$. The mortality rate, used modeling the cost of cooperation was 0 for non-cooperators and variable (into each experiment series) for cooperators. For all simulations the “fitness differential” was 0.6. The “Fitness differential” is the difference between the predation rate of isolated individuals and that for cooperators being part of a group of cooperators.

For each one of the spatial structures studied (Random Graphs[3], Small-World Networks[19] and Scale Free Networks[1]) we considered two modes of horizontal information transmission: one using fixed rates (using three different rates: $pT \in \{0, 0.5, 1\}$) and one where transmission rate is proportional to the group size of neighbors of different kind. The three variations of fixed rates plus the proportional rate gave us four scenarios of horizontal transmission.

In all scenarios production of new agents for was uniform (50/50), and the rate pT determined the probability an agent would imitate the behavior (cooperate or not) of the majority of its neighbors.

Our experiments consisted in 4 series of simulations, corresponding to the described scenarios, over a specific spatial structure. In each series (21 simulations) we varied the cost of cooperation (mortality rate of cooperators) in steps of 0.05, to cover the interval between 0 and 1. Each experiment was run 20 times to get the average of data.

For each network we set its parameters to get the same mean degree. We choose mean degree 4 in order to make a “fair” comparison between networks and with 2-D grids.



Each network structure possess a set of statistical properties that allow explain their behavior. Even though significant properties are still been developed, the most studied are: the mean geodesic path, the clustering coefficient, the degree distribution, the network resilience, the mixing patterns, the degree correlations and the community structure[10].

In table 1 we present the values of some of these properties for studied networks. Degree related values were taken from our networks, while the average path length for complex networks was calculated using equations form [10]. We also checked that degree distributions corresponds to theoretical predictions.



4. RESULTS

The results of our experiments are shown in Figure 1, where each sub-figure summarizes the results from simulations of a particular complex network structure: Random Graphs (Fig. 1(a)), Small World Networks (Fig. 1(b)) and Scale Free Networks (figure 1(c)). These figures show the final proportion of cooperators for each scenario under various cooperation cost. In all the spatial structures and all the scenarios the proportion of cooperators decreases monotonically as cooperation costs (cc) increment. Also, cooperators are the majority of the population in most circumstances.



In Random Graphs (Fig. 1(a)) curves for $pT = 0.5$ and $pT = 0$ have the same shape that $pT = 0$, but there is a



(a) Random Graph. Erdős-Renyi Model ($G(10000, 0.0002)$); parameters: $size = 10000$, $edge_probability = 0.0002$.



(b) Small World Graph. Newman-Watts model ($G(10000, 2, 0.01)$); parameters: $size = 10000$, $connections_by_direction = 2$, $rewiring_probability = 0.01$.



(c) Scale Free Networks. Barabasi-Albert model ($G(10000,2)$); parameters: $size = 10000$, $m = 2$.

Figure 1: Final fraction of cooperators in simulations with different costs for cooperation (x axis) and with different kinds of horizontal transmission of information (pT). On the left side of each sub-figure we present a graphical representation of the subjacent spatial structure.

Graph	Parameters	mean degree	degree' SD	average path length
Random Graph	G(10000,2)	3.99	1.99	6.64
Small World Graph	G(10000,2,0.01)	4	0	1250
Scale Free Network	G(10000, 4)	3.99	6.88	30615

Table 1: Grids and network properties' values.

difference of at most 15% between $pT = 0$ and $pT = 1$, being $pT = 1$ below $pT = 0$. The $pT = 0.5$ curve have separation from the $pT = 0$ curve is less than a 3%.

For Small World Graphs (Fig. 1(b)) all curves have the same shape, but $pT = 1$ is up to 50% below the $pT = 0$ curve; and the $pT = 0.5$ curve up to 15% below the $pT = 0$ curve. This network shows the biggest differences between the different information transmission rates.

The Scale Free Networks curves follow the same shapes than the 2-D grids, with smaller differences between pT 's values.

Is interesting to note that when $pT = 0$, the curves show no differences between all the spatial structures studied; and that when $pT = 0.5$ is very similar to $pT = \text{proportional}$. Large values of pT ($pT = 1$) provide the largest reduction of cooperators, especially in Small World Graphs.

5. DISCUSSION

The first feature that strikes us is that horizontal transmission of information has a negative effect on the amount of cooperators that survive the evolutionary dynamics unless the cost for cooperation are very high. This result is due to the fact that the cooperative strategy is more susceptible to invasion by the opposite strategy than the non-cooperative one. A lack of communication or horizontal transmission of information favors groups of cooperators, benefiting from each others neighborhood, who are then less likely to be disrupted by non-cooperators

The results show that some spatial networks but not all affect the effect of information transmission on cooperation dynamics. The largest effect was seen in Small World Graphs which happen to be the networks with the lowest variance in the degree of connectivity of the nodes. The effect on the evolutionary dynamics of Random Graphs and Scale Free Networks was similar. This result makes us wonder what characteristic of Small World Graph produces this difference. As with the result discussed above, a more homogeneous network makes it less likely that isolated groups of cooperators benefit from each other neighborhood. Thus the low variance in connectivity of Small World Graphs makes them less prone to nurture groups of cooperators.

These results confirm that the spatial structure affects the cooperation dynamics under horizontal information transmission. Despite this susceptibility, however, it is curious to note that the type of transmission of information modeled did not seem to affect the outcome. That is, simulations with $pT = 0.5$ were undistinguishable from those with $pT = \text{proportional}$. This indicates that flexibility, or the lack of it, in implementing the majority rule has no effect on the evolutionary dynamics of cooperation. This result unveils an additional resilience for the evolution of cooperation.

6. REFERENCES

- [1] A.-L. Barabási and R. Albert. Emergence of scaling in random networks. *Science*, 286:509–512, 1999.
- [2] R. Cipriani and K. Jaffe. On the dynamics of grouping. In *Proceedings of the Fifth IASTED International Conference on Modelling, Simulation and Optimization*, pages 56–60. Acta Press, 2005.
- [3] P. Erdős and A. Rényi. On random graphs. *Publicationes Mathematicae*, 6:290–297, 1959.
- [4] W. Hamilton. The genetic evolution of social behaviour. papers i and ii. *Journal of Theoretical Biology*, 7:1–16, 17–52, 1964.
- [5] W. Hamilton. Geometry for the selfish herd. *Journal of Theoretical Biology*, 31:295–311, 1971.
- [6] P. Hammerstein, editor. *Genetic and cultural evolution of cooperation*. Dahlem Workshops Reports. MIT Press, 2003.
- [7] C. Hauert. Fundamental clusters in spatial 2x2 games. *Proc. R. Soc. Lond. B*, 268:761–769, 2001.
- [8] K. Jaffe and R. Cipriani. Culture outsmarts nature in the evolution of cooperation. Available from <http://jasss.soc.surrey.ac.uk/10/1/7.html>, 2006.
- [9] R. Kurzban and D. Houser. Experiments investigating cooperative types in humans: A complement to evolutionary theory and simulations. *PNAS*, 102:1803–1807, 2005.
- [10] M. E. J. "M. Newman. "the structure and function of complex networks". *SIAM Review*, 45(2):167–256, 2003.
- [11] R. Nelson and S. Winter. *Evolutionary theory of economic change*. Belknap Press, 1982.
- [12] M. Nowak and K. Sigmund. The dynamics of indirect reciprocity. *Journal of Theoretical Biology*, 195:561–574, 1998.
- [13] M. Nowak and K. Sigmund. Evolutionary dynamics of biological games. *Science*, 303:793–799, 2004.
- [14] M. Nowak and K. Sigmund. Evolution of indirect reciprocity. *Nature*, 437:1291–1298, 2005.
- [15] M. A. Nowak and R. M. May. Evolutionary games and spatial chaos. *Nature*, 359:826–829, Oct. 1992.
- [16] H. Ohtsuki, C. Hauert, E. Lieberman, and M. A. Nowak. A simple rule for the evolution of cooperation on graphs and social networks. *Nature*, 441:502–505, 2006.
- [17] P. Richardson, J. Strassmann, and C. Hughes. *Not by Genes Alone: How Culture Transformed Human Evolution*. Chicago Univ. Press, 2004.
- [18] R. Trivers. The evolution of reciprocal altruism. *Quarterly Review of Biology*, 46:35–57, 1971.
- [19] D. Watts and S. Strogatz. Collective dynamics of 'small-world' networks. *Nature*, 393:440–442, 1998.